

Appendix C. Source and Accuracy of Estimates

SOURCE OF DATA

The SIPP universe is the noninstitutionalized resident population living in the United States. This population includes persons living in group quarters, such as dormitories, rooming houses, and religious group dwellings. Crew members of merchant vessels, Armed Forces personnel living in military barracks, and institutionalized persons, such as correctional facility inmates and nursing home residents, were not eligible to be in the survey. Also, U.S. citizens residing abroad were not eligible to be in the survey. Foreign visitors who work or attend school in this country and their families were eligible; all others were not eligible. With the exceptions noted above, persons who were at least 15 years of age at the time of the interview were eligible to be interviewed in the survey.

The 1985 panel SIPP sample is located in 230 primary sampling units (PSU's) each consisting of a county or a group of contiguous counties. Within these PSUs, expected clusters of two to four living quarters (LQ's) were systematically selected from lists of addresses prepared for the 1980 decennial census to form the bulk of the sample. To account for LQ's built within each of the sample areas after the 1980 census, a sample was drawn of permits issued for construction of residential LQ's up until shortly before the beginning of the panel.

In jurisdictions that do not issue building permits, small land areas were sampled and the LQ's within were listed by field personnel and then subsampled. In addition, sample LQ's were selected from a supplemental frame that included LQs identified as missed in the 1980 census and group quarters.

The first interview was conducted during February, March, April, and May of 1985. Approximately one-fourth of the sample was interviewed in each of these months. Each sample person was visited every 4 months thereafter. At each interview, the reference period was the 4 months preceding the interview month.

Approximately 17,800 living quarters were originally designated for the 1985 sample. At the first interview, interviews were obtained from the occupants of about 13,400 of the 17,800 designated LQ's for the 1985 panel. Most of the remaining 4,400 LQ's in the 1985 panel were found to be vacant, demolished, converted to nonresidential use, or otherwise ineligible for the survey. However, approximately 1,000 of the 4,400 LQ's for the 1985 panel were not interviewed because the

occupants refused to be interviewed, could not be found at home, were temporarily absent, or were otherwise unavailable. Thus, occupants of about 93 percent of all eligible living quarters participated in the first interview of the survey.

For subsequent interviews, only original sample persons (those interviewed in the first interview) and persons living with them were eligible to be interviewed. Original sample persons were followed if they moved to a new address, unless the new address was more than 100 miles from a SIPP sample area. Then, telephone interviews were attempted. All first interview noninterviewed households were automatically designated as noninterviews for all subsequent interviews. When original sample persons moved to remote parts of the country, moved without leaving a forwarding address or refused to be interviewed, additional noninterviews resulted.

A person was classified as interviewed or noninterviewed for the entire 1985 panel based on the following definitions. Interviewed sample persons were defined to be 1) those for whom self or proxy responses were obtained for each reference month of all eight interviews or 2) those for whom self or proxy responses were obtained for the first reference month of the panel and for each subsequent reference month until they were known to have died or moved to an ineligible address (foreign living quarters, institutions, or military barracks). Noninterviewed persons were defined to be those for whom neither self nor proxy responses were obtained for one or more reference months of the eight interviews (but not because they were deceased or moved to an ineligible address).

For longitudinal estimates, approximately 36,500 persons were counted as initially interviewed in the 1985 panel. In the 1985 panel weighting procedure, approximately 23,000 persons were classified as interviewed. Persons who missed interviews due to the February 1986 sample cut were not classified as noninterviews but were adjusted for in the weighting procedure by a special factor. The person nonresponse rate is estimated to be 37 percent for the entire 1985 panel. Some respondents did not respond to some of the questions; therefore, the overall nonresponse rate for some items, especially sensitive income and money related items, is higher than the person nonresponse rate.

ESTIMATION

Several stages of weight adjustments were involved in the estimation procedure used to derive the SIPP longitudinal person weights. Each person received a

base weight equal to the inverse of his/her probability of selection. Two noninterview adjustment factors were applied. One adjusted the weights of interviewed persons in interviewed households to account for households which were eligible for the sample but could not be interviewed at the first interview. The second was applied to compensate for person noninterviews occurring in subsequent interviews. The Bureau has used complex techniques to adjust the weights for nonresponse, but the success of these techniques in avoiding bias is unknown. Another factor was applied to each interviewed person's weight to account for the SIPP sample areas not having the same population distribution as the strata from which they were selected.

An additional stage of adjustment to longitudinal person weights was performed to reduce the mean square error of the survey estimates. This was accomplished by bringing the sample estimates into agreement with monthly Current Population Survey (CPS) type estimates of the civilian (and some military) noninstitutional population of the United States by demographic characteristics including age, sex, race, and Hispanic ethnicity as of the specified control date. The CPS estimates by age, race, sex, and Hispanic origin were themselves brought into agreement with estimates from the 1980 decennial census which have been adjusted to reflect births, deaths, immigration, emigration, and changes in the Armed Forces since 1980.

In weighting of the 1985 panel, two first stage factors were incorrect. The impact of this error on primary SIPP estimates is believed to be minimal.

The data for the longitudinal estimates provided in the report are obtained from all eight interviews of the 1985 panel of the Survey of Income and Program Participation (SIPP).

ACCURACY OF ESTIMATES

SIPP estimates are based on a sample; they may differ somewhat from the figures that would have been obtained if a complete census had been taken using the same questionnaire, instructions, and enumerators. There are two types of errors possible in an estimate based on a sample survey: nonsampling and sampling. We are able to provide estimates of the magnitude of SIPP sampling error, but this is not true of nonsampling error. Found in the next sections are descriptions of sources of SIPP nonsampling error, followed by a discussion of sampling error, its estimation, and its use in data analysis.

Nonsampling variability. Nonsampling errors can be attributed to many sources, e.g., inability to obtain information about all cases in the sample, definitional difficulties, differences in the interpretation of questions, inability or unwillingness on the part of the respondents

to provide correct information, inability to recall information, errors made in collection such as in recording or coding the data, errors made in processing the data, errors made in estimating values for missing data, biases resulting from the differing recall periods caused by the interviewing pattern used, and failure of all units in the universe to have some probability of being selected for the sample (undercoverage). Quality control and edit procedures were used to reduce errors made by respondents, coders and interviewers.

Undercoverage in SIPP results from missed living quarters and missed persons within sample households. It is known that undercoverage varies with age, race, and sex. Generally, undercoverage is larger for males than for females and larger for Blacks than for non-Blacks. Ratio estimation to independent age-race-sex population controls partially corrects for the bias due to survey undercoverage. However, biases exist in the estimates to the extent that persons in missed households or missed persons in interviewed households have characteristics different from those of interviewed persons in the same age-race-sex group. Further, the independent population controls used have not been adjusted for undercoverage.

Comparability with other estimates. Caution should be exercised when comparing data from this report with data from other SIPP publications or with data from other surveys. The comparability problems are caused by such sources as the seasonal patterns for many characteristics, different nonsampling errors, and different concepts and procedures.

Sampling variability. Standard errors indicate the magnitude of the sampling error. They also partially measure the effect of some nonsampling errors in response and enumeration, but do not measure any systematic biases in the data. The standard errors for the most part measure the variations that occurred by chance because a sample rather than the entire population was surveyed.

USES AND COMPUTATION OF STANDARD ERRORS

Confidence intervals. The sample estimate and its standard error enable one to construct confidence intervals, ranges that would include the average result of all possible samples with a known probability. For example, if all possible samples were selected, each of these being surveyed under essentially the same conditions and using the same sample design, and if an estimate and its standard error were calculated from each sample, then:

1. Approximately 68 percent of the intervals from one standard error below the estimate to one standard error above the estimate would include the average result of all possible samples.

2. Approximately 90 percent of the intervals from 1.6 standard errors below the estimate to 1.6 standard errors above the estimate would include the average result of all possible samples.
3. Approximately 95 percent of the intervals from two standard errors below the estimate to two standard errors above the estimate would include the average result of all possible samples.

The average estimate derived from all possible samples is or is not contained in any particular computed interval. However, for a particular sample, one can say with a specified confidence that the average estimate derived from all possible samples is included in the confidence interval.

Hypothesis testing. Standard errors may also be used for hypothesis testing, a procedure for distinguishing between population characteristics using sample estimates. The most common types of hypotheses tested are 1) the population characteristics are identical versus 2) they are different. Tests may be performed at various levels of significance, where a level of significance is the probability of concluding that the characteristics are different when, in fact, they are identical.

All statements of comparison in the report have passed a hypothesis test at the 0.10 level of significance or better. This means that, for differences cited in the report, the estimated absolute difference between parameters is greater than 1.6 times the standard error of the difference.

To perform the most common test, compute the difference $X_A - X_B$, where X_A and X_B are sample estimates of the characteristics of interest. A later section explains how to derive an estimate of the standard error of the difference $X_A - X_B$. Let that standard error be s_{DIFF} . If $X_A - X_B$ is between -1.6 times s_{DIFF} and $+1.6$ times s_{DIFF} , no conclusion about the characteristics is justified at the 10 percent significance level. If, on the other hand, $X_A - X_B$ is smaller than -1.6 times s_{DIFF} or larger than $+1.6$ times s_{DIFF} , the observed difference is significant at the 10 percent level. In this event, it is commonly accepted practice to say that the characteristics are different. Of course, sometimes this conclusion will be wrong. When the characteristics are, in fact, the same, there is a 10 percent chance of concluding that they are different.

Note that as more tests are performed, more erroneous significant differences will occur. For example, at the 10 percent significance level, if 100 independent hypothesis tests are performed in which there are no real differences, it is likely that about 10 erroneous differences will occur. Therefore, the significance of any single test should be interpreted cautiously.

Note concerning small estimates and small differences. Summary measures are shown in the report only when the base is 200,000 or greater. Because of the

large standard errors involved, there is little chance that estimates will reveal useful information when computed on a base smaller than 200,000. Also, nonsampling error in one or more of the small number of cases providing the estimate can cause large relative error in that particular estimate. Estimated numbers are shown, however, even though the relative standard errors of these numbers are larger than those for the corresponding percentages. These smaller estimates are provided primarily to permit such combinations of the categories as serve each user's needs. Therefore, care must be taken in the interpretation of small differences since even a small amount of nonsampling error can cause a borderline difference to appear significant or not, thus distorting a seemingly valid hypothesis test.

Standard error parameters and tables and their use.

Most SIPP estimates have greater standard errors than those obtained through a simple random sample because clusters of living quarters are sampled for the SIPP. To derive standard errors that would be applicable to a wide variety of estimates and could be prepared at a moderate cost, a number of approximations were required. Estimates with similar standard error behavior were grouped together and two parameters (denoted "a" and "b") were developed to approximate the standard error behavior of each group of estimates. Because the actual standard error behavior was not identical for all estimates within a group, the standard errors computed from these parameters provide an indication of the order of magnitude of the standard error for any specific estimate. These "a" and "b" parameters vary by characteristic and by demographic subgroup to which the estimate applies. Table C-1 provides base "a" and "b" parameters to be used for 1985 longitudinal panel estimates.

For those users who wish further simplification, we have also provided general standard errors in tables C-2 and C-3. Note that these standard errors must be adjusted by a factor from table C-1. The standard errors resulting from this simplified approach are less accurate. Methods for using these parameters and tables for computation of standard errors are given in the following sections.

Standard errors of estimated numbers. The approximate standard error, s_x , of an estimated number of persons shown in this report can be obtained in two ways.

It may be obtained by the use of the formula

$$s_x = fs \quad (1)$$

where f is the appropriate "f" factor from table C-1 and s is the standard error of the estimate obtained by interpolation from table C-2. Alternatively, s_x may be approximated by the formula

$$s_x = \sqrt{ax^2 + bx} \quad (2)$$

Table C-1. **SIPP Generalized Variance Parameters for Estimates Using Panel Weights: 1985 Longitudinal Panel File**

Characteristic	a	b	f factor
TOTAL OR WHITE PERSONS			
16+ program participation and benefits, and poverty (3):			
Both sexes.....	-0.0001965	33,468	.90
Male.....	-0.0004134	33,468	.90
Female.....	-0.0003743	33,468	.90
16+ income and labor force (4):			
Both sexes.....	-0.0000669	11,410	.52
Male.....	-0.0001411	11,410	.52
Female.....	-0.0001276	11,410	.52
All others ¹ (5):			
Both sexes.....	-0.0001800	41,497	1.00
Male.....	-0.0003722	41,497	1.00
Female.....	-0.0003485	41,497	1.00
BLACK PERSONS			
Poverty (1):			
Both Sexes.....	-0.0010347	28,548	.83
Male.....	-0.0022234	28,548	.83
Female.....	-0.0019352	28,548	.83
All others (2):			
Both Sexes.....	-0.0005564	15,351	.61
Male.....	-0.0011956	15,351	.61
Female.....	-0.0010406	15,351	.61

¹These parameters are to be used for all tabulations not specifically covered by any other category in this table, e.g., for retirement and pension tabulations, for 0+ benefits, 0+ poverty, 0+ income, and 0+ labor force tabulations.

Note:For cross-tabulations, apply the parameters of the category showing the smaller number in parentheses.

Here x is the estimated number and "a" and "b" are the parameters associated with the particular type of characteristic. Use of formula (2) will provide more accurate results than the use of formula (1).

Illustration. Suppose that we have a SIPP estimate of 960,000 adults 18 years or over who were poor in 1985 yet exited poverty in 1986 and increased their weeks or hours worked. The appropriate "a" and "b" parameters to use in calculating a standard error for the estimate are obtained from table C-1. They are a = -.0001965 and b = 33,468, respectively. Using formula (2), the approximate standard error is

$$\sqrt{(-.0001965) (960,000)^2 + (33,468) (960,000)} = 179,000$$

The 90-percent confidence interval is from 674,000 to 1,246,000. Therefore, a conclusion that the average estimate derived from all possible samples lies within a range computed in this way would be correct for roughly 90 percent of all samples.

Using formula (1), the appropriate "f" factor (f = .90) from table C-1, and the appropriate standard error of the estimate from table C-2, the approximate standard error is

$$s_x = .90 (198,680) = 179,000$$

Standard errors of estimated percentages. This section refers to the percentages of a group of persons possessing a particular attribute such as the percentage of persons who were poor in the two-year reference

period. The reliability of an estimated percentage, computed using sample data for both numerator and denominator, depends upon both the size of the percentage and the size of the total upon which the percentage is based. Estimated percentages are relatively more reliable than the corresponding estimates of the numerators of the percentages, particularly if the percentages are over 50 percent. For example, the percent of persons in poverty is more reliable than the estimated number of persons in poverty. When the numerator and denominator of the percentage have different parameters, use the parameter (and appropriate factor) of the

Table C-2. **Standard Errors of Estimated Numbers of Persons for 1985 Longitudinal Panel File**

(Numbers in thousands)

Size of estimate	Standard error	Size of estimate	Standard error
200.....	91	22,000.....	909
300.....	112	26,000.....	978
600.....	158	30,000.....	1041
1,000.....	203	50,000.....	1275
2,000.....	287	80,000.....	1472
5,000.....	451	100,000.....	1533
8,000.....	566	130,000.....	1534
11,000.....	659	150,000.....	1475
13,000.....	713	200,000.....	1049
15,000.....	763	220,000.....	646
17,000.....	808	230,000.....	149

numerator. If proportions are presented instead of percentages, note that the standard error of a proportion is equal to the standard error of the corresponding percentage divided by 100.

For the percentage of persons, the approximate standard error, $s_{(x,p)}$, of the estimated percentage, p , can be obtained by the formula

$$s_{(x,p)} = fs \tag{3}$$

where f is the appropriate “ f ” factor from table C-1, and s is the standard error of the estimate obtained by interpolation from table C-3. Alternatively, it may be approximated by the formula

$$s_{(x,p)} = \sqrt{\frac{b}{x} p(100-p)} \tag{4}$$

Here x is the base of the percentage, p is the percentage ($0 > p > 100$) and b is the “ b ” parameter associated with the characteristic in the numerator. Use of this formula will give more accurate results than use of formula (3).

Illustration. Suppose that 28 percent of the 3,682,000 adults who exited poverty in 1986 did not work in either 1985 or 1986. Using formula (4) and the “ b ” parameter of 33,468 from table C-1, the approximate standard error is

$$\sqrt{\frac{33,468}{3,682,000} (28)(100-28)} = 4.3 \text{ percent}$$

Consequently, the 90 percent confidence interval is from 21.1 to 34.9 percent.

Standard error of a difference. The standard error of a difference between two sample estimates, x and y , is equal to

$$s_{(x-y)} = \sqrt{s_x^2 + s_y^2} \tag{5}$$

where s_x and s_y are the standard errors of the estimates x and y . The estimates can be numbers, averages, percents, ratios, etc. The above formula assumes that the correlation coefficient, r , between the characteristics estimated by x and y is zero. If r is really positive (negative), then this assumption will result in a tendency towards overestimates (underestimates) of the true standard error.

Illustration. Suppose that, 23.9 percent of the adults who exited poverty in 1986 increased their weeks or hours worked, whereas, 28 percent of such adults did not work in either 1985 or 1986. Using the appropriate b parameter from table C-1 and formula (4), the standard errors of these percentages are approximately 4.1 percent and 4.3 percent, respectively.

The standard error of the difference is computed using formula (5):

$$\sqrt{(4.1)^2 + (4.3)^2} = 5.9 \text{ percent}$$

Suppose that it is desired to test at the 10 percent significance level whether the above two percentages differ significantly. To perform the test, compare the

Table C-3. Standard Errors of Estimated Percentages of Persons for 1985 Longitudinal Panel File

Base of estimated percentage (thousands)	Estimated percentage					
	1 or 99	2 or 98	5 or 95	10 or 90	25 or 75	50
200	4.5	6.4	9.9	13.7	19.7	22.8
300	3.7	5.2	8.1	11.2	16.1	18.6
600	2.6	3.7	5.7	7.9	11.4	13.1
1,000	2.0	2.9	4.4	6.1	8.8	10.2
2,000	1.4	2.0	3.1	4.3	6.2	7.2
5,000	0.9	1.3	2.0	2.7	3.9	4.6
8,000	0.7	1.0	1.6	2.2	3.1	3.6
11,000	0.6	0.9	1.3	1.8	2.7	3.1
13,000	0.6	0.8	1.2	1.7	2.4	2.8
17,000	0.5	0.7	1.1	1.5	2.1	2.5
22,000	0.4	0.6	0.9	1.3	1.9	2.2
26,000	0.4	0.6	0.9	1.2	1.7	2.0
30,000	0.4	0.5	0.8	1.1	1.6	1.9
50,000	0.3	0.4	0.6	0.9	1.2	1.4
80,000	0.2	0.3	0.5	0.7	1.0	1.1
100,000	0.2	0.3	0.4	0.6	0.9	1.0
130,000	0.2	0.3	0.4	0.5	0.8	0.9
180,000	0.2	0.2	0.3	0.5	0.7	0.8
200,000	0.1	0.2	0.3	0.4	0.6	0.7
220,000	0.1	0.2	0.3	0.4	0.6	0.7
230,000	0.1	0.2	0.3	0.4	0.6	0.7

numerator. If proportions are presented instead of percentages, note that the standard error of a proportion is equal to the standard error of the corresponding percentage divided by 100.

For the percentage of persons, the approximate standard error, $s_{(x,p)}$, of the estimated percentage, p , can be obtained by the formula

$$s_{(x,p)} = fs \tag{3}$$

where f is the appropriate “ f ” factor from table C-1, and s is the standard error of the estimate obtained by interpolation from table C-3. Alternatively, it may be approximated by the formula

$$s_{(x,p)} = \sqrt{\frac{b}{x} p(100-p)} \tag{4}$$

Here x is the base of the percentage, p is the percentage ($0 > p > 100$) and b is the “ b ” parameter associated with the characteristic in the numerator. Use of this formula will give more accurate results than use of formula (3).

Illustration. Suppose that 28 percent of the 3,682,000 adults who exited poverty in 1986 did not work in either 1985 or 1986. Using formula (4) and the “ b ” parameter of 33,468 from table C-1, the approximate standard error is

$$\sqrt{\frac{33,468}{3,682,000} (28)(100-28)} = 4.3 \text{ percent}$$

Consequently, the 90 percent confidence interval is from 21.1 to 34.9 percent.

Standard error of a difference. The standard error of a difference between two sample estimates, x and y , is equal to

$$s_{(x-y)} = \sqrt{s_x^2 + s_y^2} \tag{5}$$

where s_x and s_y are the standard errors of the estimates x and y . The estimates can be numbers, averages, percents, ratios, etc. The above formula assumes that the correlation coefficient, r , between the characteristics estimated by x and y is zero. If r is really positive (negative), then this assumption will result in a tendency towards overestimates (underestimates) of the true standard error.

Illustration. Suppose that, 23.9 percent of the adults who exited poverty in 1986 increased their weeks or hours worked, whereas, 28 percent of such adults did not work in either 1985 or 1986. Using the appropriate b parameter from table C-1 and formula (4), the standard errors of these percentages are approximately 4.1 percent and 4.3 percent, respectively.

The standard error of the difference is computed using formula (5):

$$\sqrt{(4.1)^2 + (4.3)^2} = 5.9 \text{ percent}$$

Suppose that it is desired to test at the 10 percent significance level whether the above two percentages differ significantly. To perform the test, compare the

Table C-3. Standard Errors of Estimated Percentages of Persons for 1985 Longitudinal Panel File

Base of estimated percentage (thousands)	Estimated percentage					
	1 or 99	2 or 98	5 or 95	10 or 90	25 or 75	50
200	4.5	6.4	9.9	13.7	19.7	22.8
300	3.7	5.2	8.1	11.2	16.1	18.6
600	2.6	3.7	5.7	7.9	11.4	13.1
1,000	2.0	2.9	4.4	6.1	8.8	10.2
2,000	1.4	2.0	3.1	4.3	6.2	7.2
5,000	0.9	1.3	2.0	2.7	3.9	4.6
8,000	0.7	1.0	1.6	2.2	3.1	3.6
11,000	0.6	0.9	1.3	1.8	2.7	3.1
13,000	0.6	0.8	1.2	1.7	2.4	2.8
17,000	0.5	0.7	1.1	1.5	2.1	2.5
22,000	0.4	0.6	0.9	1.3	1.9	2.2
26,000	0.4	0.6	0.9	1.2	1.7	2.0
30,000	0.4	0.5	0.8	1.1	1.6	1.9
50,000	0.3	0.4	0.6	0.9	1.2	1.4
80,000	0.2	0.3	0.5	0.7	1.0	1.1
100,000	0.2	0.3	0.4	0.6	0.9	1.0
130,000	0.2	0.3	0.4	0.5	0.8	0.9
180,000	0.2	0.2	0.3	0.5	0.7	0.8
200,000	0.1	0.2	0.3	0.4	0.6	0.7
220,000	0.1	0.2	0.3	0.4	0.6	0.7
230,000	0.1	0.2	0.3	0.4	0.6	0.7

difference of 4.1 percent to the product of $1.6 \times 5.9 = 9.4$ percent. Since the percent difference is smaller than 1.6 times the standard error of the difference, the data do not support the hypothesis that the two percent estimates are significantly different at the 10 percent level.

Standard errors of ratios of means and medians.

The standard error for a ratio of means or medians is approximated by:

$$s_{\frac{x}{y}} = \sqrt{\left(\frac{x}{y}\right)^2 \left[\left(\frac{s_y}{y}\right)^2 + \left(\frac{s_x}{x}\right)^2 \right]} \quad (6)$$

where x and y are means or medians, and s_x and s_y are their associated standard errors. Formula 6 assumes that the means or medians are not correlated. If the correlation between the population means or medians

estimated by x and y are actually positive (negative), then this procedure will tend to produce overestimates (underestimates) of the true standard error for the ratio of means or medians.

Illustration. Suppose the SIPP estimate of Black-to-White median family income per person in 1986 is .62. Also, suppose that the median family income per person and its corresponding standard error are \$17,822 and \$669, respectively, for the Blacks; and \$28,962 and \$474, respectively, for the Whites.

Using formula (6), the standard error for this ratio is approximated by:

$$\frac{s_x}{y} = \sqrt{\left(\frac{17,822}{28,962}\right)^2 \left[\left(\frac{669}{17,822}\right)^2 + \left(\frac{474}{28,962}\right)^2 \right]} = .025$$